

Recurrent rhabdomyolysis and glutaric aciduria type I: a case report and literature review

Gu-Ling Qian, Fang Hong, Fan Tong, Hai-Dong Fu, Ai-Min Liu

Hangzhou, China

Background: Glutaric acidemia type I (GA-I) is a rare metabolic disorder caused by mutation of the glutaryl-CoA dehydrogenase (*GCDH*) gene. The occurrence of rhabdomyolysis with GA-I is extremely rare.

Methods: We reported a child with recurrent rhabdomyolysis and undiagnosed glutaric acidemia type I (GA-I). And a literature review was performed.

Results: A 4.5-year-old girl was admitted to our hospital due to recurrent rhabdomyolysis for 3 times within three years. At the third admission, she was diagnosed with GA-I by biochemical testing and mutation analysis. The girl was found to have a serine to leucine replacement mutation of the *GCDH* gene in exon 8 at position 764. Other three patients with rhabdomyolysis and GA-I were discovered by literature searching.

Conclusion: This report highlights that patients with GA-I may have an increased risk of rhabdomyolysis.

World J Pediatr 2016;12(3):368-371

Key words: *GCDH* mutation; glutaric acidemia type I; rhabdomyolysis

Introduction

Glutaric acidemia type I (GA-I), a rare (1:100 000) metabolic disorder, is caused by the mutation of the glutaryl-CoA dehydrogenase (*GCDH*) gene.^[1] This mutation results in *GCDH* deficiency and the accumulation of lysine, hydroxylysine, and tryptophan and their

intermediate breakdown products including glutaric acid, glutaryl-CoA, 3-hydroxyglutaric acid, and glutaconic acid.^[1] The accumulation of 3-hydroxyglutaric acid and glutaconic acid can lead to damage to the brain and other organs. Macrocephaly at birth and later movement disorders, typically occurring between 6 and 18 months of age, are key manifestations of GA-I.^[1] High blood and/or urine glutarylcarnitine (C5DC) detected at newborn screening is suggestive of GA-I.^[2] This report describes a patient with rhabdomyolysis and GA-I. This is a noteworthy case because the occurrence of rhabdomyolysis associated with GA-I is extremely rare.

Case report

A 2.5-year-old girl was admitted to the hospital for 2-day fatigue, paleness, high fever (>39°C), and a convulsion. On physical examination, her muscle strength was measured grade 4 (Table 1) and urine was dark brown. Routine urine and blood tests revealed: 3+ urine protein and 2 red blood cells/high power field; hemoglobin (Hb), 121 g/L (reference: 110-130); alanine aminotransferase (ALT), 3050 U/L (reference: 7-35); lactate dehydrogenase (LDH), 3990 U/L (reference: 140-280); creatine kinase (CK), 108640 U/L (reference: 134-391); CK-MB, 1794 U/L (reference: 10-13); and blood myoglobin, 257.4 µg/L (reference: 0.1-70). As rhabdomyolysis was diagnosed, the girl received symptomatic treatment including fluid supplementation and urine alkalization for 3 days. Echocardiography and ecocardiography showed nothing abnormal, suggesting no myocardial damage. Her urine was clear, and muscle

Author Affiliations: Department of Inherited Metabolic Disease (Qian GL, Hong F, Tong F) and Department of Nephrology (Fu HD, Liu AM), Children's Hospital, Zhejiang University School of Medicine, Hangzhou, China

Corresponding Author: Hai-Dong Fu, MD, Department of Nephrology, Children's Hospital, Zhejiang University School of Medicine, 57 Zhuganxiang, Hangzhou 310003, China (Tel: +86-13396553987; Email: fhdhz@163.com)

doi: 10.1007/s12519-016-0042-x

©Children's Hospital, Zhejiang University School of Medicine, China and Springer-Verlag Berlin Heidelberg 2016. All rights reserved.

Table 1. Muscle strength grading

Grade 0	No muscle contraction.
Grade 1	Trace contraction is noted in the muscle by palpating while the patient attempts to contract it.
Grade 2	Able to actively move the muscle when gravity is eliminated.
Grade 3	Move the muscle against gravity, but not against resistance from the examiner.
Grade 4	Move the muscle group against some resistance from the examiner.
Grade 5	Move the muscle group and overcome the resistance of the examiner. This is the normal muscle strength.

strength returned to grade 5.

One year later, she was re-hospitalized for shortness of breath and fatigue for 3 days. Her muscle strength was grade 2/3, and urine color was normal. Urine testing showed negative urine protein and 2 red blood cells/high power field; Hb, 125 g/L; ALT, 1209 U/L; LDH, 6140 U/L; and CK, 43 440 U/L. Spinal fluid analysis revealed a leukocyte count of 2×10^6 cells/L (reference: $4-11 \times 10^6$); CK, 227 U/L (reference: 0-22); and total protein, 336.9 mg/L (reference: <450). Paneth's globulin was negative. Cranial magnetic resonance image revealed a bilateral arachnoid cyst. Muscle biopsy was performed to rule out a possible underlying defect in muscle metabolism,^[3] and pathological examination revealed decreased muscle fiber density (Fig.). The girl received urine alkalization and cell membrane stabilization for two weeks, and her muscle strength returned to normal.

At the age of 4.5 years, the girl was admitted to the hospital again because of running nose and vomiting for 3 days and difficulty in breathing for half a day. She suffered from mild respiratory distress and her muscle strength was grade 2. Her urine was yellow in color, and the results of blood tests were as follows: ALT, 2020 U/L; CK, 175 640 U/L; CK-MB, 1150 U/L; Hb, 132 g/L; and myoglobin, 509 μ g/L (reference: 0-70). Blood

analysis using Tandem mass spectrometry revealed C5DC, 1.02 mmol/mol creatinine (reference: 0.02-0.14); C5DC/C8, 17 (reference: 0.19-7); and C5DC/C16, 1.96 (reference: 0.01-0.07). She was treated by urine alkalization and cell membrane stabilization, and after three weeks her muscle strength returned to grade 4/5. Finally, she was discharged from the hospital.

Gas chromatography-mass spectrometry analysis of urine organic acids revealed: glutaric acid, 126.99 μ mol/mmol creatinine (reference: 1.9-4); and 3-hydroxyglutaric acid, 3.39 μ mol/mmol creatinine (reference: 0), consistent with GA-I. *GCDH* gene analysis on the patient, her parents, and sisters revealed a missense mutation in exon 8 (C764>T), leading to TCG>TTG and Ser>Leu. The girl had a homozygous mutation, whereas her parents and two sisters had a heterozygous mutation. DNA analysis of blood samples from 50 healthy subjects confirmed that this was a mutation, not a polymorphism.

After the diagnosis of GA-I, the patient was treated with carnitine (50 mg/kg per day), a protein/amino acid supplement (60 g/day) and a predominantly vegetarian and protein diet. Biochemical testing at the age of 5.1 years revealed no abnormalities; at the age of 5 years and 3 months, C5DC was 1.08, and glutaric acid and 3-hydroxyglutaric acid were 19.21 μ mol/mmol




Fig. A: Quadriceps biopsy revealed decreased muscle fiber density (hematoxylin and eosin staining, original magnification $\times 400$); **B:** Axial magnetic resonance image showing a bilateral arachnoid cyst in the middle cranial fossa (arrows) on T2 image.

Table 2. Summary of rhabdomyolysis in patients with glutaric aciduria type I (GA I)

Case	GA I			Rhabdomyolysis			
	Age/sex	History/age at confirmation	Treatments	Recurrent	Cause	Clinical manifestations	Outcome/follow-up
Wilson (1999)	4 y/M	Yes/NA	Carnitine and riboflavin supplement	Yes	Viral infection or unknown	Fever, shock, tachypnea, encephalopathy, myoglobinuria	Death
Chow (2003)	4.5 y/F	Yes/after birth	Protein restriction, carnitine and riboflavin supplement	No	Viral infection	Fever, shock, encephalopathy, myoglobinuria	Death
Jamuar (2012)	8 y/M	Yes/4 months	Low protein diet, carnitine and riboflavin supplement	Yes	Febrile illness	Fever, respiratory distress, dystonia, encephalopathy	Recurrent dystonia and rhabdomyolysis
Present	5 y/F	No	No standard treatment before diagnosis	Yes	Unknown	Fever, fatigue, respiratory distress, convulsion, myoglobinuria	Stable without symptoms

F: female; M: male; NA: not available. *: The patient experienced two episodes of mild rhabdomyolysis, and was severely disabled at 13 years of age.

creatinine and 5.22 $\mu\text{mol}/\text{mmol}$ creatinine, respectively. The patient had normal growth and development without symptoms during a 3.5 year of follow-up.

Discussion

By searching Medline, the Cochrane Library, EMBASE, and Google Scholar, we found three patients with rhabdomyolysis and GA-I (Table 2).^[4-6] These patients aged from 4 to 8 years (except for our patient) suffered from GA-I, which was treated with carnitine and riboflavin, and protein was restricted. Recurrent rhabdomyolysis was thought to be caused by viral infection in two patients^[4,5] and febrile illness in one.^[6] Fever as a symptom of rhabdomyolysis occurred in all patients, and other common symptoms included encephalopathy, shock, myoglobinuria, and respiratory distress. Of the three patients, 2 died, and one became disabled.

Our patient had a serine to leucine replacement mutation of the *GCDH* gene in exon 8 at position 764. To our knowledge, this is a novel *GCDH* gene mutation. Findings of a study indicate that a single mutation (IVS10-2A>C) is the predominant one in patients with GA-I.^[7] Overall, there are more than 200 *GCDH* gene mutations,^[8,9] including R402W missense mutation.^[11] Other *GCDH* mutations in Chinese with GA-I include A219T, R386G, IVS3+1G>A, G178R, and R355H,^[10] c.245G>C (p.Arg82Pro),^[11] and missense mutations (c.406G>T, C.416C>G, c.442G>A, c.640A>G, c.901G>A, c.979G>A, c.1207C>T), frameshift mutations (c.873delC, c.1172-1173insT, c.1282-1285ins71), and nonsense mutations (c.411C>G).^[12] *GCDH* mutations include c.281G>A (p.Arg94Gln), c.401A>G (p.Asp134Gly), c.662T>C (p.Leu221Pro), c.881G>C (p.Arg294Pro), c.1173dupG (p.Asn392Glufs*5), c.1238A>G (p.Tyr413Cys) and c.1241A>C (p.Glu414Ala) in Indian patients;^[13] 89 or 90delC, Y155C, IVS4+2T>C, G244S, Q352X, G354A, K361E, 1144-1145delGC, and S305L in Japanese;^[14] and G390V and X439W in Italians and Portuguese.^[15]

In our patient and those reported elsewhere, the most obvious potential precipitating factor for rhabdomyolysis is altered bioenergetics due to the metabolic disorder caused by *GCDH* deficiency. Indeed, genetic inborn errors of metabolism are known to cause rhabdomyolysis.^[16] Investigation of *GCDH* in mice showed markedly decreased brain and skeletal muscle CK activity after administration of lysine.^[17] Before each attack, our patient consumed a large amount of mutton which contains large amount of lysine.

Glutaryl-CoA dehydrogenase enzyme activity of fibroblasts or leukocytes is the "gold standard" for the diagnosis of rhabdomyolysis.^[18] However, Kölker et

al^[18] reported that 2 disease-causing mutations and elevated 3-OH-GA and GA in urine were sufficient to diagnose the disease. Our patient had elevated 3-OH-GA and GA in urine (by gas chromatography-mass spectrometry) and a homozygous mutation (missense mutation in exon 8, C764>T); her parents and 2 sisters had a heterozygous mutation. These findings are sufficient to diagnose GA-I.

In summary, we described the fourth patient with rhabdomyolysis and GA-I. Compared with the reported patients, our patient had neither a history of GA-I or obvious manifestations of the condition nor any adverse neurological outcomes. We believe that the present case may help to identify risk factors and causes of rhabdomyolysis in patients with GA-I.

Funding: This study was supported by the National Natural Science Foundation of China (81170664).

Ethical approval: This study was approved by the Ethical Committee, Children's Hospital, Zhejiang University School of Medicine. This case report was prepared in accordance with the Health Insurance Portability and Accountability Act (HIPAA) regulations. The patient's parents/legal guardians provided informed consent for data collection and publication.

Competing interest: The authors have no conflict of interest related to the treatment of the patient or reporting.

Contributors: Qian GL guaranteed the integrity of the entire study, and prepared the manuscript. Hong F performed the literature research. Tong F contributed to the study concepts. Fu HD was responsible for the manuscript editing. Liu AM contributed to the manuscript review.

References

- Hedlund GL, Longo N, Pasquali M. Glutaric acidemia type 1. *Am J Med Genet C Semin Med Genet* 2006;142C:86-94.
- Couce ML, López-Suárez O, Bóveda MD, Castiñeiras DE, Cocho JA, García-Villoria J, et al. Glutaric aciduria type I: outcome of patients with early- versus late-diagnosis. *Eur J Paediatr Neurol* 2013;17:383-389.
- Bosch X, Poch E, Grau JM. Rhabdomyolysis and acute kidney injury. *N Engl J Med* 2009;361:62-72.
- Wilson CJ, Collins JE, Leonard JV. Recurrent rhabdomyolysis in a child with glutaric aciduria type I. *J Inher Metab Dis* 1999;22:663-664.
- Chow SL, Rohan C, Morris AA. Rhabdomyolysis in glutaric aciduria type I. *J Inher Metab Dis* 2003;26:711-712.
- Jamuar SS, Newton SA, Prabhu SP, Hecht L, Costas KC, Wessel AE, et al. Rhabdomyolysis, acute renal failure, and cardiac arrest secondary to status dystonicus in a child with glutaric aciduria type I. *Mol Genet Metab* 2012;106:488-490.
- Tang NL, Hui J, Law LK, To KF, Mak TW, Cheung KL, et al. Overview of common inherited metabolic diseases in a Southern Chinese population of Hong Kong. *Clin Chim Acta* 2001;313:195-201.
- Goodman SI, Stein DE, Schlesinger S, Christensen E, Schwartz M, Greenberg CR, et al. Glutaryl-CoA dehydrogenase mutations

- in glutaric acidemia (type I): review and report of thirty novel mutations. *Hum Mutat* 1998;12:141-144.
- 9 Zschocke J, Quak E, Guldborg P, Hoffmann GF. Mutation analysis in glutaric aciduria type I. *J Med Genet* 2000;37:177-181.
- 10 Tang NL, Hui J, Law LK, Lam YY, Chan KY, Yeung WL, et al. Recurrent and novel mutations of *GCDH* gene in Chinese glutaric acidemia type I families. *Hum Mutat* 2000;16:446.
- 11 Liu Q, Chen Y, Chen W. Mutation analysis of *GCDH* gene in four patients with glutaric acidemia type I. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2015;32:187-191. [In Chinese]
- 12 Wang Q, Li X, Ding Y, Liu Y, Song J, Yang Y. Clinical and mutational spectra of 23 Chinese patients with glutaric aciduria type I. *Brain Dev* 2014;36:813-822.
- 13 Gupta N, Singh PK, Kumar M, Shastri S, Gulati S, Kumar A, et al. Glutaric acidemia type I-clinico-molecular profile and novel mutations in *GCDH* gene in Indian patients. *JIMD Rep* 2015 2015;21:45-55.
- 14 Mushimoto Y, Fukuda S, Hasegawa Y, Kobayashi H, Purevsuren J, Li H, et al. Clinical and molecular investigation of 19 Japanese cases of glutaric acidemia type I. *Mol Genet Metab* 2011;102:343-348.
- 15 Busquets C, Soriano M, de Almeida IT, Garavaglia B, Rimoldi M, Rivera I, et al. Mutation analysis of the *GCDH* gene in Italian and Portuguese patients with glutaric aciduria type I. *Mol Genet Metab* 2000;71:535-537.
- 16 Zhang RN, Li YF, Qiu WJ, Ye J, Han LS, Zhang HW, et al. Clinical features and mutations in seven Chinese patients with very long chain acyl-CoA dehydrogenase deficiency. *World J Pediatr* 2014;10:119-125.
- 17 Amaral AU, Cecatto C, Seminotti B, Zanatta Â, Fernandes CG, Busanello EN, et al. Marked reduction of Na(+), K(+)-ATPase and creatine kinase activities induced by acute lysine administration in glutaryl-CoA dehydrogenase deficient mice. *Mol Genet Metab* 2012;107:81-86.
- 18 Kölker S, Christensen E, Leonard JV, Greenberg CR, Boneh A, Burlina AB, et al. Diagnosis and management of glutaric aciduria type I--revised recommendations. *J Inher Metab Dis* 2011;34:677-694.

Received April 9, 2015

Accepted after revision August 5, 2015